

The client sample was analyzed for plant-based cannabinoids by Liquid Chromatography (LC). The collected data was compared to data collected for certified reference standards at known concentrations.

47450-CN

ID	Weight %	Conc.			
D9-THC	ND	ND			
THCV	ND	ND			
CBD	1.88 wt %	17.92 mg/mL			
CBDV	0.01 wt %	0.06 mg/mL			
CBG	0.01 wt %	0.06 mg/mL			
CBC	0.07 wt %	0.71 mg/mL			
CBN	0.01 wt %	0.07 mg/mL			
THCA	ND	ND			
CBDA	ND	ND			
CBGA	ND	ND			
D8-THC	ND	ND			
exo-THC	ND	ND			
Total	1.98 wt%	18.82 mg/mL	0%	Cannabinoids (wt%)	1.9%
Max THC	-	-			
Max CBD	1.88 wt%	17.92 mg/mL			

Max THC (and Max CBD) are calculated values for total cannabinoids after heating, assuming complete decarboxylation of the acid to the neutral form. It is calculated based on the weight loss of the acid group during decarboxylation: Max THC = $(0.877 \times THCA) + THC$. This calculation does not include other cannabinoid isomers (eg. D8-THC and exo-THC). ND = None detected above the limits of detection (LLD)

420 Fortune Blvd • Milford, MA 01757 • 617-221-3356 www.ProVerdeLabs.com

The client sample was analyzed by Head-Space Gas Chromatography (HS-GC). The collected data was compared to data collected for certified reference standards at known concentrations.

⁴⁷⁴⁵⁰⁻TP

Compound	wt%	Quantitative Profile	Compound	wt%	Quantitative Profile		
			Humulene				
Sabinene*	0.013		P-cymene				
Menthol*			Camphene				
A-phellandrene*			B-pinene	0.107			
Myrcene			Eucalyptol				
Isopulegol			A-terpenine				
Nerolidol-cis			3-carene				
G-terpenine	0.094		A-pinene	0.014			
Nerolidol-trans			Limonene	0.693			
A-bisabolol			Geraniol				
Linalool			Ocimene-2				
B-caryophyllene			Ocimene-1				
Caryophyllene Oxide			Terpinolene				
Guaiol							
Wi	t% 0.00	0.50 1.	00	0.0	0 0.50 1.00		
Total Terpene: 0.9 wt%							

* Indicates semi-qualitative calculation based on recorded peak areas.

END OF REPORT